



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

2893 [1921, 184]. Proposed by **NATHAN ALTSHILLER-COURT**, University of Oklahoma.

Find the locus of the mid-point of the segment determined by two given skew lines in a variable plane turning about a fixed axis, not coplanar with either of the given lines.

SOLUTION BY **J. K. WHITTEMORE**, Yale University.

Let the axis of the variable plane be the z axis, and suppose the x and y axes chosen parallel one to each of the given skew lines. This coördinate system is generally oblique. The equations of the skew lines are (1) $y = b_1, z = c_1$; (2) $x = a_2, z = c_2$; the variable plane, $y = \lambda x$, intersects the two lines in points of coördinates, (1) $x = b_1/\lambda, y = b_1, z = c_1$; (2) $x = a_2, y = \lambda a_2, z = c_2$. The coördinates of the mid-point of the segment, a point of the required locus, are

$$x = \frac{a_2}{2} + \frac{b_1}{2\lambda}, \quad y = \frac{b_1}{2} + \frac{\lambda a_2}{2}, \quad z = \frac{c_1 + c_2}{2}.$$

Eliminating the parameter λ , the equations of the required locus are

$$\left(x - \frac{a_2}{2}\right)\left(y - \frac{b_1}{2}\right) = \frac{a_2 b_1}{4}, \quad z = \frac{c_1 + c_2}{2}.$$

The locus is a hyperbola lying in a plane parallel to both given skew lines and half way between them; its asymptotes are parallel to the given skew lines; its center is the point of intersection of the diagonals of the parallelepiped, three of whose edges lie on the axis of the variable plane and the given skew lines; it intersects the axis of the variable plane.

Also solved by **WILLIAM HOOVER**.

2900 [1921, 277]. Proposed by **I. A. BARNETT**, University of Saskatchewan.

AB is the diameter of a circle and Q_0 any point on the circumference; Q_1, Q_2, Q_3, \dots , are the points of bisection of the arcs AQ_0, AQ_1, AQ_2, \dots ; to prove that the product of the chords of the circle $BQ_1, BQ_2, BQ_3, \dots, BQ_n$ is equal to $OA^n \cdot (AQ_0/AQ_n)$, O being the center of the circle.

SOLUTION BY **A. M. HARDING**, University of Arkansas.

Arc $AQ_1 = \text{arc } Q_1Q_0$. Hence, $\angle OBQ_1 = \angle AQ_0Q_1 = \angle Q_1AQ_0$. Hence, the isosceles triangles, OBQ_1 and Q_1Q_0A are similar. Then,

$$\frac{BQ_1}{AQ_0} = \frac{OB}{AQ_1}; \quad \text{that is,} \quad BQ_1 = OA \cdot \frac{AQ_0}{AQ_1}.$$

In a similar manner, it may be shown that

$$\begin{aligned} BQ_2 &= OA \cdot \frac{AQ_1}{AQ_2}, \\ BQ_3 &= OA \cdot \frac{AQ_2}{AQ_3}, \\ &\vdots \\ BQ_n &= OA \cdot \frac{AQ_{n-1}}{AQ_n}. \end{aligned}$$

Multiplying these equations gives

$$BQ_1 \cdot BQ_2 \cdot BQ_3 \cdot \dots \cdot BQ_n = OA^n \cdot \frac{AQ_0}{AQ_n}.$$

Also solved by **T. M. BLAKSLEE**, **ARTHUR PELLETIER**, and **A. V. RICHARDSON**.

2945 [1922, 29]. Proposed by **T. M. BLAKSLEE**, Ames, Iowa.

A point P in the plane of the triangle ABC rotates in a given direction around the vertices taken in either cyclical order, in each case through an angle equal to the corresponding angle of the triangle. That is, for example, AP rotates around A through an angle equal to the angle A of the triangle; then BP around B through an angle equal to the angle B , and so on. Prove